

All Databases PubMed Nucleotide Protein Genome Structure OMIM PMC Journals Books
 Search PubMed for [Advanced Search](#)

[Limits](#) [Preview/Index](#) [History](#) [Clipboard](#) [Details](#)

Display Show Sort By Send to

All: 1 Review: 0

1: [Biomaterials](#). 2005 Jun;26(17):3503-9.

ELSEVIER [Links](#)
FULL-TEXT ARTICLE

Effect of laser therapy on attachment, proliferation and differentiation of human osteoblast-like cells cultured on titanium implant material.

[Khadra M](#), [Lynqstadaas SP](#), [Haanaes HR](#), [Mustafa K](#).

Department of Oral Surgery and Oral Medicine, Faculty of Dentistry, University of Oslo, P.O.Box 1109 Blindern, N-0317 Oslo, Norway. maawan@odont.uio.no

The aim of this in vitro study was to investigate the effect of low-level laser therapy (LLLT) on the attachment, proliferation, differentiation and production of transforming growth factor- β 1 (TGF- β 1) by human osteoblast-like cells (HOB). Cells derived from human mandibular bone were exposed to GaAlAs diode laser at dosages of 1.5 or 3 J/cm² and then seeded onto titanium discs. Non-irradiated cultures served as controls. After 1, 3 and 24h, cells were stained and the attached cells were counted under a light microscope. In order to investigate the effect of LLLT on cell proliferation after 48, 72 and 96 h, cells were cultured on titanium specimens for 24h and then exposed to laser irradiation for three consecutive days. Specific alkaline phosphatase activity and the ability of the cells to synthesize osteocalcin after 10 days were investigated using p-nitrophenylphosphate as a substrate and the ELSA-OST-NAT immunoradiometric kit, respectively. Cellular production of TGF- β 1 was measured by an enzyme-linked immunosorbent assay (ELISA), using commercially available kits. LLLT significantly enhanced cellular attachment ($P < 0.05$). Greater cell proliferation in the irradiated groups was observed first after 96 h. Osteocalcin synthesis and TGF- β 1 production were significantly greater ($P < 0.05$) on the samples exposed to 3 J/cm². However, alkaline phosphatase activity did not differ significantly among the three groups. These results showed that in response to LLLT, HOB cultured on titanium implant material had a tendency towards increased cellular attachment, proliferation, differentiation and production of TGF- β 1, indicating that in vitro LLLT can modulate the activity of cells and tissues surrounding implant material.

PMID: 15621240 [PubMed - indexed for MEDLINE]

Display Show Sort By Send to

Related articles

The effect of low level laser irradiation on implant-tissue interaction. In vivo [Swed Dent J Suppl. 2005]

Determining optimal surface roughness of TiO₂ blasted titanium implant [Clin Oral Implants Res. 2001]

Determining optimal dose of laser therapy for attachment and pro [J Biomed Mater Res A. 2005]

Review Underlying mechanisms at the bone-surface interface during [J Periodontol Res. 1997]

Review Implant surface characteristics modulate differentiation behavior of [Adv Dent Res. 1999]

» See reviews... | » See all...

Cited by 1 PubMed Central article

Visible light regulates neurite outgrowth of nerve cells. [Cytotechnology. 2007]

Recent Activity

[Turn Off](#) [Clear](#)

Effect of laser therapy on attachment, proliferation and differentiation of human

The effect of low level laser irradiation on implant-tissue interaction. In vivo and in

Wound healing in cell studies and animal model experiments by Low Level Laser

Low level laser therapy (Classes I, II and III) for treating rheumatoid arthritis.

Low level laser therapy (Classes I, II and III) for treating osteoarthritis.

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Privacy Statement](#) | [Freedom of Information Act](#) | [Disclaimer](#)